

Osteoporosis Clinical Practice Guidelines

Definition

The world health organization WHO defines osteoporosis as a systemic skeletal disease characterized by low bone mass and micro -architectural deterioration of bone tissue with a resultant increase in fragility and risk of fracture

Assessment (History and Examination)

Initial Evaluation for Osteoporosis

All postmenopausal women age ≥ 50 years of age should undergo clinical assessment for osteoporosis and a detailed history, physical exam, and clinical fracture risk assessment with Fracture Risk Assessment tool (FRAX®) Note: FRAX age 40-90, Saudi FRAX in process of endorsement. Use USA white as per Saudi Osteoporosis Society SOS1).Or the Kuwaiti FRAX (similar hip fracture incidence to Saudi Arabia) until the Saudi FRAX is available

| History | Exam |
|---|---|
| <ul style="list-style-type: none"> •Prior osteoporosis-related fractures •Prolonged steroid use •Height loss > 6 cm historically •Current smoking •Excess alcohol ≥ 3 units per day •Parental hip fracture •Falls in past 12 months •Other high-risk conditions or medications | <ul style="list-style-type: none"> •Height loss (>2cm prospectively) •Weight (BMI) •Low <60 Kg •Major loss ($\geq 10\%$ of weight since age 25) •Kyphosis •Rib to pelvis distance >2 FBs •Balance and gait, “Get up and Go” Test |
| FRAX Clinical risk factors in FRAX® www.shef.ac.uk/FRAX | |
| <ul style="list-style-type: none"> •age •Sex •body mass index (BMI) •smoking, alcohol use •prior fracture •parental history of hip fracture •use of glucocorticoids •rheumatoid arthritis •secondary osteoporosis femoral neck BMD | |



Indications for BMD Testing (1)

Menopausal women, and men aged 50-64 years

Routine Screening indicated by age

Menopausal women, and men aged 50-64 years with clinical risk factors for fracture:

- All women ≥ 40 years who have sustained low-trauma fragility fracture
- Previous fragility fracture or maternal history of hip fracture .
- hypogonadism or premature menopause (< 45 years), Prolonged secondary amenorrhea (>1 year)
- Prolonged glucocorticoid use ≥ 3 months cumulative use in the past year of prednisone-equivalent dose ≥ 7.5 mg daily
- Other high-risk medication use (tamoxifen , thiazolidinedione , Empagliflozin, PPI and anticonvulsant)
- X-ray findings suggestive of osteoporosis such as vertebral fracture, osteopenia identified on X-ray, fragility fracture, loss of height, or thoracic kyphosis (clinical or radiological finding)
- Current smoking
- High alcohol intake
- Low body weight (< 60 kg) or major weight loss ($>10\%$ of weight at age 25 years)
- Rheumatoid arthritis
- Other disorders strongly associated with osteoporosis such as primary hyperparathyroidism, type 1 diabetes, osteogenesis imperfecta, uncontrolled hyperthyroidism, Cushing's, disease, chronic malnutrition or malabsorption, chronic inflammatory conditions (e.g., inflammatory bowel disease)

All women age ≥ 60 years in Saudi Arabia (expert opinion screen)
All men age ≥ 65 years

2020 AACE American Association of Clinical Endocrinologists Diagnosis of Osteoporosis in Postmenopausal Women (2)

1. T-score -2.5 or below in the lumbar spine, femoral neck, total proximal femur, or 1/3 radius
2. Low-trauma **spine or hip fracture** (regardless of bone mineral density)
3. T-score between -1.0 and -2.5 and a fragility fracture of **proximal humerus, pelvis, or distal forearm**
4. T-score between -1.0 and -2.5 and **high FRAX[®]** (or if available, TBS-adjusted FRAX[®]) 10-year probability for major osteoporotic fracture is $\geq 20\%$ or the 10-year probability of hip fracture is $\geq 3\%$ FRAX[®] = fracture risk assessment tool; TBS = trabecular bone score


For All Osteoporotic Patient Evaluate for Causes of Secondary Osteoporosis Before Start Treatment

- Serum chemistry: TSH calcium, phosphate, total protein, albumin, liver enzymes, alkaline phosphatases, creatinine, and electrolytes.
- Serum 25-hydroxyvitamin D Complete blood cell count
- SPE serum protein electrophoresis if vertebral fracture or suspect multiple myeloma
- X-ray lateral thoracolumbar screen for vertebral fracture if not available from DXA

Recommend pharmacologic therapy Education on lifestyle measures, fall prevention, benefits, & risks of medications



Management





| | |
|---|--|
| CALCIUM Premenopausal, men <50 years and pregnant women (1000 mg/d), Postmenopausal, men >50 years (1500 mg/d) | Instructions: Calcium should be taken with meals for better absorption. Calcium should not be taken with iron (absorption may be adversely affected when given concurrently) Caution in patient with hypercalcemia and patients with history of renal stones |
| VITAMIN D Premenopausal, men <50 yr and pregnant women (600 IU/d) Postmenopausal, men >50 yr (1000 IU/d) | Instructions: Expose to sun for 10-15 min 2-3 times/wk Caution in patient with hypercalcemia and patients with history of renal stones |
| ALENDRONATE 70 mg once weekly Antiresorptive oral Consider drug holiday after 5 years | Instructions: should be taken as soon as patient wakes up in the morning, before eating, or drinking anything. - Tablet should be swallowed as a whole with a large glass (8 ounces) of plain water only (not mineral water, coffee, juice, or any other liquid). - Patient should not lie down on their back, eat, or drink for at least 30 min after taking alendronate. SE: 1) Hypocalcemia 2) atypical fracture of the femur 3) osteonecrosis of the jaw defer initiation or hold if invasive dental procedures Contraindicated: 1) Should not be prescribed for patients with active esophageal abnormalities or peptic ulcer disease. And inability to remain upright for at least ½ hour after the dose 2) In pregnancy, women who plan to be pregnant. 3) In patients with creatinine clearance below 30 mL/min. One infusion per year over minimum of 15 min. Good hydration before receiving the medication. |
| zoledronate 5 mg IV once yearly Antiresorptive intravenous Consider drug holiday after 3 years | Correct hypocalcemia before starting treatment. SE: Hypersensitivity, flu-like reaction, Risk of atypical fracture, ONJ, Arterial fibrillation Contraindicated: in pregnancy, women who plan to be pregnant, and in patients with creatinine clearance below 30 mL/min |
| Denosumab (Prolia) 60 mg SC every 6 months Antiresorptive SC injection | 60 mg denosumab in 1 mL solution in a single-use pre-filled syringe or vial Subcutaneous injection every 6 months SE: Eczema, cellulitis, low calcium Contraindicated: 1) In pregnancy, women who plan to be pregnant. 2) Risk of atypical fracture, ONJ, Dose adjustment for renal impairment is not necessary. CrCl ≥ 30 mL/min risk of severe hypocalcemia if CrCl <30 mL/min |
| RALOXIFEN 60 mg/d oral SERMS (Evista) | Stop in periods of prolonged immobilization (surgery, long flight, cholestyramine intake) SE: premenopausal women worsening of hot flashes, leg cramps, increase risk of deep vein thrombosis. Increased risk of thromboembolic events Risks needs to be weighed against benefits, especially in patients with or at risk of CHD (in whom treatment reduces vertebral fracture and breast cancer risk at the same absolute rate that it increases the VTE and fatal stroke risk) |



Recommend: Education on lifestyle measures, fall prevention, benefits and risks of medications

| Exercise type /benefits | Frequency | Comment |
|--|----------------------|---|
| Posture exercises keep you standing tall, not stooped. | Daily 10 mint | Pay attention to your posture posture when you stand and sit, do back exercises that extend your spine. |
| Balance exercises help you be more stable on your feet. You can walk more easily. Good balance helps prevent falls. | Daily 20 mint | walk heel to toe, reduce base of support, shift your weight, respond to things that upset your balance. |
| Strength exercises keep you strong and fit. | 2 times per week | Exercise for leg ,arm ,chest shoulder and back. Use body weight against gravity ,band and weights * |
| Aerobic physical activity (moderate to vigorous intensity) improves your overall health. It can reduce your risk of disease. It may improve your bone strength. | 150 minutes per week | Do aerobic physical activity for about 20 to 30 minutes per day. Exercise for at least 10 minutes at a time. In total, do 150 minutes or more per week.* If you are new to exercise or if you have had a spine fracture, start at low to moderate intensity — 3 to 6 on the scale* |

*Refer to physical therapy for advice for proper exercise for each patient

Give patient medication card when starting the treatment this is essential for collaborative medical care between specialist and primary care example: Abaloparatide or Teriparatide taken once in life time for up to 2 years and need to be followed by antiresorptive treatment. Moreover, it is essential to know when the patient can go for drug holiday.

Medication: _____

Calcium: Dietary sources: _____ mg Supplements: _____ mg

Vitamin D:


Exercise: _____ minutes daily / weekly

Fall Prevention advice _____

Follow up DXA / labs in _____ months. Return visit in _____ months



R3 Osteoporosis Screening Pathway:



| APPROVAL | | | |
|---------------------------|---|----------------|------------|
| | Name: | Position: | Signature: |
| Prepared By: | Dr. Ahmed Al Zahrani Dr. Ahlam Al Harbi | FM Consultants | |
| Reviewed and Approved By: | Dr. Mansoor Allajhar Dr. Musa Althwayee Dr. Ahmed Al Zahrani Dr. Hajar Al Suma Dr. Ahlam Al Harbi | FM Consultants | |

Reference

- 2015 Guidelines for Osteoporosis in Saudi Arabia: Recommendations from the Saudi Osteoporosis Society Ann Saudi Med 2015;35(1):1-12
1. AMERICAN ASSOCIATION OF CLINICAL ENDOCRINOLOGISTS/ AMERICAN COLLEGE OF ENDOCRINOLOGY CLINICAL PRACTICE GUIDELINES FOR THE DIAGNOSIS AND TREATMENT OF POSTMENOPAUSAL OSTEOPOROSIS - 2020 UPDATE Endor Pract. 2020;26(Suppl 1)
 2. Camacho et al. Endocr Pract. 2020.26.564-570
 3. Shoback et al. J Clin Endocrinol Metab, 2020, 105(3), 1-8
 4. Eastell et al. J Clin Endocrinol Metab. 2019, 104, 1595-1622
 5. Kanis et al. Osteoporosis Int. 2020, 31, 1-12

